

FOLIAR FEED II

FOLIAR APPLICATION

Cotton: First application of 1-2 qt. per acre at 4 leaf stage, second when cotton begins to square. Repeat at 10-14 day intervals.

Field Crops: 3 applications of 1-2 qts. per acre in enough water to assure good coverage. Apply at or before bloom and repeat at 10-14 day intervals.

Tree Crops: After fruit set, apply 2-3 qts. per 100 gallons of water. May be repeated at 10-14 day intervals.

Turf Grass, Hay Meadows, and Bermuda grass pastures: Apply 2-4 qts. with enough water to assure good coverage. (Additional rates with increased volumes of water.) Apply when grass begins to green and then at 20-30 day intervals.

STORAGE INSTRUCTIONS

FOLIAR FEED II is a specifically formulated acid based fertilizer. It is corrosive to brass and mild steel. Bulk FFII must be stored in stainless steel, fiberglass, or polyethylene storage tanks. Fittings and valves must be PVC plastic or stainless only! Pail storage should be limited to a maximum of 2 pallets high with each pallet being 2 pails high.

A FOLIAR Plant Food containing available Nitrogen, Phosphoric Acid, Potash, Boron, Chelated iron, Chelated Zinc, Chelated Manganese, and Chelated Copper

GUARANTEED ANALYSIS

Total Nitrogen (N)	15.00%
Available Phosphoric Acid (P ₂ O ₅)	10.00%
Soluble Potash (K ₂ O).....	5.00%
Boron (B)02%
Chelated Copper (Cu)05%
Chelated Iron (Fe)10%
Chelated Manganese (Mn).....	.05%
Chelated Zinc (Zn).....	.05%

Nutrients derived from Urea, Orthophosphoric acid, muriate of potash, zinc sulfate, manganese sulfate, ferrous sulfate, boric acid, and copper sulfate. Also contains ethylene diamine tetraacetic acid (EDTA) and other chelating agents.

-CAUTION-

Corrosive liquid. Keep out of reach of children. Causes eye and skin irritation. May be harmful if swallowed. In case of contact flush with water for at least 15 minutes and get medical help.



THE J.C. SMITH CO.
SAN SABA, TEXAS 76877

• 2.5 gallon	25 lb	11.36 kg
• 5 gallon	50 lb	22.73 kg
• 275 gallon	2750 lb	1250 kg

PREPLANT SOIL APPLICATION

Broadcast: Apply 5 gal. per acre in enough water for even distribution.

Row Application: Apply 3 gal. per acre in enough water for even distribution. Apply 2 inches to the side and 2 inches below seed row.

Plant deficiency problems usually occur during the early stages of growth. Foliar Feed II should be applied at or before this time for maximum efficiency. Foliar Feed II tends to stimulate root growth, and to assure essential nutrients at the critical growth stages of fruiting and flowering plants. Foliar Feed II is compatible with most fungicides and insecticides, and acts as a buffering agent to lower PH in the spray tank and to help control hydrolysis of insecticides and fungicides. Foliar Feed II goes with good management practices to give better yields and more dollars per acre return for your investment.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability of fitness for a particular purpose expressed or implied, extends to the use of this product contrary to label instruction or under abnormal conditions, or under conditions not reasonably foreseeable to the seller and buyer assumes the risk of any such use.