

Pasture Booster 32-10XTR

Liquid Feed Supplement

Bulk Liquid Feed
For Beef Cattle on Pasture

GUARANTEED ANALYSIS

| | |
|--|--------------|
| Crude Protein, min..... | 32.00% |
| <i>(This includes not more than 19.50% equivalent crude protein from non-protein nitrogen)</i> | |
| Crude, Fiber, max..... | 0.10% |
| Crude Fat, min..... | 10.00% |
| Calcium, min..... | 0.25% |
| Calcium, max..... | 1.00% |
| Phosphorus, min..... | 0.95% |
| Potassium, min..... | 0.90% |
| Magnesium (Mg), min..... | 0.18% |
| Cobalt (Co), min..... | 1ppm |
| Copper (Cu), min..... | 25ppm |
| Iron (Fe), min..... | 150ppm |
| Manganese (Mn), min..... | 60ppm |
| Zinc (Zn), min..... | 80ppm |
| Selenium (Se), min..... | .3ppm |
| Iodine (I), min..... | 2.5ppm |
| Vitamin A, min..... | 15,000 IU/lb |
| Vitamin D, min..... | 2,500 IU/lb |
| Vitamin E, min..... | 10 IU/lb |
| Moisture, max..... | 55.00% |

INGREDIENTS: Condensed Corn Extractives, Ammonium polyphosphate solution, Processed polyunsaturated vegetable oil refined lipid, Urea, Condensed extracted glutamic Acid fermentation product, zinc sulfate, manganese sulfate, ferrous sulfate, copper sulfate, ethylene diaminedihydriodide, cobalt carbonate, sodium selenite, Vitamin A Propionate, D-activated Animal Sterol (source of Vitamin D3), dl-alpha tocopheryl acetate (source of Vitamin E).

*****WARNING*****

THIS FEED SUPPLEMENT SHOULD BE USED ONLY IN ACCORDANCE WITH DIRECTION FURNISHED ON THIS LABEL.

Feeding Directions

1. Feed to ruminants only.
2. Provide Unlimited drinking water.
3. Unlimited roughage must be available free choice. Overgrazed pastures are not adequate.
4. Grain may be fed as desired.
Provide unlimited amounts of the following separately free choice: Salt (loose or block), ground limestone and /or an adequate mineral supplement
5. For cattle on pasture feed free choice in covered lick tanks and feed 1-6 lbs per head per day. For cattle fed total mixed rations include from 1-6 lbs. per head per day. Pasture Booster 32-10 should be used at a level not to exceed 10% of total daily dry matter feed intake. Consumption should be monitored to obtain desired daily intake. Additional care should be exercised with starved, stressed or debilitated animals.
6. Damaged tanks, broken or open tops must be repaired or replaced before cattle are allowed access.
This product must be agitated before being dispensed or fed.

For additional information, consult your dealer.

Seller makes no warranty, expressed or implied, as to description, utility, productiveness or any other matter, except as to guaranteed analysis.

**Manufactured by:
The J.C. Smith Co.
San Saba, Texas 76877**